

Renormalization Group Approach to the Problem of Flow Through Irregular Packed Beds

Dmitri Volchenkov and Ricardo Lima

CNRS, Centre de Physique Theorique, 13288 Marseille Cedex 09, Luminy Case 907, France

volchen@cptsg2.univ-mrs.fr, lima@cptsg2.univ-mrs.fr

February 1, 2008

Abstract

We have calculated the asymptotics of Green's function of the differential equation of nonlinear diffusion in the microscopic range with strong porosity fluctuations in the problem of flow through irregular packed beds for the arbitrary dimension of space and arbitrary porosity fluctuations covariance.

PACS numbers: 47.55 M, 11.10.G, 47.10. +g, 05.40. +j, 42.27. Gs

1 Introduction

Properties of flows in porous media are of great interest of modern industry and technology. In spite of practical importance and long-standing attention paid to this problem there is still no solidarity on model of the flow through the packed beds.

One can point out an approach of a single particle diffusing with random dynamics in a Poissel velocity field, which goes back to the classical work of Taylor [1]. Nevertheless, the experimental measurements are still described by the usual diffusion equation where a modified dispersion coefficient is used; see [2] and references therein. The problem of these considerations is of the essentially gaussian solutions of the diffusion equation. The numerous experimental observations, [2] demonstrates clearly that the previous approach needs to be supplemented.

One of the first attempts to overcome this limitation is proposed in [3], where a Markov random process is used to model the diffusion of impurities in a turbulent stream. In [4] the stochastic field was introduced to model the porosity instead of the diffusion process itself.

Their result for the one-dimensional Fick law of diffusion for some marker transported by a carrier fluid through an irregularly but statistically homogeneous packed column leads to the appearance in the equation of the mean concentration of a time diffusion term. Together with a source term they are to be responsible for the departure of the solutions from gaussian.

In this paper we keep on study the diffusion equation of the Fick-type in which the porosity ε fluctuates strongly along the column axis. There are many methods of defining porosity; see [5] and references therein. Each of them replaces the complex network of voids with a single number that represents an average property. In the present work we use porosity as a point quantity in a porous medium and define an elementary volume to characterize a medium macroscopically following [6] and [7]. The size of the elementary volume around a point P we suppose to be smaller than the total medium so that it can represent a fluid flow at P . We suppose also that there is enough pores to allow statistical averaging. Since porosity varies, the maximum length is the characteristic length ξ that indicates the rate of change of porosity, and the minimum length w is the pore size.

The porosity is directly related to the size of the pores relative to the matrix. Considering the problem of packed column, we denote by x the axial coordinate of the column, $0 \leq x \leq L$, then the fraction of the column that is voids defines the averaging value of ε , $0 \leq \varepsilon \leq 1$. If P be a point inside a porous medium on the column axis, see Fig.1., surrounded by a layer of volume $V(l_i)$, one can define a ratio

$$\varepsilon_i = \frac{V_v(l_i)}{V(l_i)}$$

in which $V_v(l_i)$ is the volume of void space within volume $V(l_i)$.

Consider a sequence of values of l_i such that $l_1 > l_2 > l_3 > \dots$. For large values of $V(l_i)$ the ratio ε_i may change gradually as $V(l_i)$ gets smaller. As l falls below a certain value ξ there will be large fluctuations in porosity when $V(l)$ is approaching the dimension of the pores (this $V(\xi)$ may be taken as the elementary volume). Below this value the average value of porosity has no meaning, then it is natural to consider the porosity as a function of x , where x be a distance apart from the point P .

Flow phenomena for $x > \xi$ is well investigated in the porous medium theory; see for example [5] for a review. Below ξ , for $w \leq x < \xi$ flow properties are subject to strong microscopic effects and usually left beyond the investigations due to an extremely large fluctuations in porosity when x is approaching the dimension of the pores. However, namely this interval of scales is of great interest for the variety of problems in chemical engineering and environmental studies.

We continue to study the microscopic properties of flows through a porous media

with strongly fluctuative porosity using the quantum field theory renormalization group method (RG) and $4 - \epsilon$ expansion [8]. The technique which is developed in what following is similar to one of [9] devoted to waves propagation in a randomly inhomogeneous medium with strongly developed fluctuations.

In Sec. 2, the problem is stated and formulated in the quantum-field formalism. In Sec. 3, the Green's function for the equation of nonlinear diffusion is constructed for the case of gaussian distributed fluctuative porosity. We have introduced the effective action functional which is equivalent to the problem discussed under the certain conditions in Subsec. 3.1. Investigating the properties of the theory, we have justified the result of [4] on the inclusion of a "time-diffusive" term into the equation of nonlinear diffusion in case of fluctuating porosity for arbitrary dimension of space and for arbitrary stochastic process originating the porosity fluctuations.

In Subsec. 3.3 and Subsec. 3.4, the procedure of ultraviolet renormalization of the theory with the "time-diffusive" term included is considered and critical dimensions including anomalies are calculated. The results obtained for static correlation functions meet the well-known empirical laws of Richardson and Kolmogorov. It is shown that for the parameters taking values within the physical domain (positive viscosity, *etc.*), the renormalization-group equations have a manifold of infrared (IR) attractive fixed points. This implies the presence of a universal scaling regime in the IR domain of the microscopical region of fluctuative porosity for some range of values of physical parameters.

The main result of the present paper is the detailed description of the effect of strongly fluctuating porosity on the form of static (*i.e.*, of equal time) as well as of dynamic Green's functions of the differential equation of nonlinear diffusion.

Similar to the problems of nonlinear diffusion and of chemically active scalar admixture, the problem of the flow through the irregular packed beds with strongly fluctuative porosity, in the general case, is of an infinite number of coupling constants, *i.e.*, no one of the statistical momenta of higher order can be omitted from the consideration. However, in contrast with the problems mentioned any of these terms does not contain UV-divergences, so that the UV renormalization procedure as well as the critical behavior of the quantities are unsensitive to certain hypotheses on the porosity function as well as the certain statistical distribution for porosity in the microscopical range.

It turns out to be that fluctuative porosity does not affect on the universal properties of the static Green's functions, *i.e.*, it does not change the indices of empirical laws of Richardson and Kolmogorov. But porosity changes the amplitude factors of static correlation, *i.e.*, the Kolmogorov constant and the effective diffusivity coefficient.

Fluctuative porosity does affect the dynamic Green's functions providing the damping

retarded time-spectrum.

Qualitatively, for the concentration one has an outgoing damping traveling wave type solution which decay rate as well as an amplitude depend on porosity.

2 Formulation of the Problem

We shall consider the diffusion process of mixture in a randomly inhomogeneous packed beds with strongly fluctuative porosity in the microscopic range $w \leq |\mathbf{x}| \ll \xi$ for the arbitrary value of space dimension d .

The problem is posed as follows: To find the retarded Green's function (propagator), averaged over the imposed statistics of fluctuative porosity $\varepsilon(\mathbf{x})$ and flow velocity $\mathbf{v}(\mathbf{x}, \mathbf{t})$, of the inhomogeneous diffusion equation for the concentration $u(\mathbf{x}, \mathbf{t})$ of some marker in the available fraction of the volume:

$$\varepsilon(\mathbf{x})\partial_{\mathbf{t}}\mathbf{u}(\mathbf{x}, \mathbf{t}) + (\mathbf{v}\partial)\mathbf{u}(\mathbf{x}, \mathbf{t}) + \nu\Delta\mathbf{u}(\mathbf{x}, \mathbf{t}) = \mathbf{0} \quad (1)$$

in which ν is the Fick's diffusion coefficient, Δ is the Laplace operator, $\mathbf{v}(\mathbf{x}, \mathbf{t})$ is the velocity of fluid flow. The porosity $\varepsilon(\mathbf{x}) = \varepsilon_0 + \tilde{\varepsilon}(\mathbf{x})$ is the sum of constant and random components. From the physical point of view the value of porosity belongs to the interval $\varepsilon \in [0, 1]$, so that, generally speaking, $\tilde{\varepsilon}$ cannot be taken as a gaussian distributed function. To model porosity one can use any function which values lie within the interval $[0, 1]$; for a qualitative determinacy in what following we model it by the function

$$\varepsilon(\mathbf{x}) = (\mathbf{1} + \sin \mathbf{c}^{-1}\varphi(\mathbf{x}))/\mathbf{2}, \quad (2)$$

where $c \equiv \nu/\xi$ is a parameter with the dimensionality of velocity which characterizes the velocity of the diffusion process in the scale ξ , and $\varphi(\mathbf{x})$ is a gaussian distributed phase function (of the same dimensionality) with zero mean value and static correlation function of the Ornstein-Zernike type, [10]

$$D_\varphi \equiv \langle \varphi(\mathbf{x})\varphi(\mathbf{y}) \rangle = \eta_0 \frac{\nu_0^3}{\mathbf{c}_0^2} \frac{1}{(2\pi)^d} \int d\mathbf{k} (\mathbf{k}^2 + \mu^2)^{(2-d-2\delta)/2} \exp i\mathbf{k}(\mathbf{x} - \mathbf{y}), \quad (3)$$

where $\mu \simeq 1/\xi$ is a mass parameter. We consider all parameters in (3) as bare (non-renormalized) parameters and supply them with the index "0". For future convenience in (3) we have introduced the amplitude factor by such a way that it meets the proper dimensionality of the correlation. The coupling constant η_0 plays the role of an expansion parameter in the perturbation theory such that the relation $\eta_0\nu_0 \equiv \Lambda^{2\delta}$ defines the characteristic maximum momentum scale in the theory, $\Lambda \simeq 1/w$.

The parameter of the regular expansion of correlation functions 2δ is the deviation of the space dimensionality from 2. We demonstrate below that under the certain physical assumptions the real value of parameter δ is $\delta_r = 1$.

The dragging velocity v along with the fluctuating component of porosity $\tilde{\varepsilon}$ prescribe a certain dimensionless parameter in the problem which is analogous to the Reynolds number, $Re_\varepsilon = \tilde{\varepsilon}vw/\nu$, where $\tilde{\varepsilon}$ is characteristic porosity fluctuations in the scale of w , and v is a characteristic fluctuative flow velocity in a pore. A statistical steady state is expected when $Re_\varepsilon \gg 1$. In terms of the energy dissipation of fluid flow the parameter Re_ε defines a dissipation length l_d by the relation $Re_\varepsilon^{4/3} \sim l_d w^{-1}$. Starting from some values Re_ε , the dissipation length falls into the range of microscopic effects, $w \ll r \ll \xi$. It leads to the formation of inertial range of scales, $l_d \ll \xi$, which is substantially analogous to the Kolmogorov's dissipation free range [16].

The basic results of the phenomenological Kolmogorov-Obukhov theory [12] can be generalized directly to the problem in question: Assume that in the inertial range of wave numbers ($1/\xi \ll k \ll 1/l_d$) correlation functions of φ depend not only on the mean energy dissipation rate W of fluid flow but also on an additional argument, the velocity of diffusion process, c . Then, assuming that the correlation functions do not depend on viscosity, for an equal-time pair correlation function of φ , one has

$$D(k) \sim W^{2/3} k^{-11/3} \mathcal{F}(W^{2/3} k^{-2/3} c^{-2}), \quad (4)$$

where $\mathcal{F}$ is a scaling function of dimensionless arguments. By the way, (4) predicts the Kolmogorov's scaling ($\Delta_\varphi = -1/3$) for the stochastic phase field φ . This phenomenological result is taken into account in the model correlation (3) by choosing $\delta_r = 1$ as a real value for δ . In accordance with (4) and [18], one obtains the scaling dimensions for quantities in the amplitude factor of (3) $\Delta_c = -1/3$ and $\Delta_\nu = -2/3$; for $\delta = 1$ (3) provides exactly the Kolmogorovian scaling dimension for the correlation $\langle \varphi \varphi \rangle$.

Similarly, to describe the effect of eddy diffusion which obviously plays an essential role in the microscopic diffusion process we consider the dragging velocity to be a d -dimensional gaussian vector field $\mathbf{v}(\mathbf{x}, t)$ of zero mean value and of covariance:

$$D_v \equiv \langle v_i(\mathbf{x}, t) v_j(\mathbf{y}, t') \rangle = g_0 \nu_0 \frac{\delta(t' - t)}{(2\pi)^d} \int d\mathbf{k} P_{ij}(k) (k^2 + m^2)^{-\frac{d}{2} - \frac{\epsilon}{2}} \exp i\mathbf{k}(\mathbf{x} - \mathbf{y}), \quad (5)$$

where $P_{ij}(k) = \delta_{ij} - k_i k_j / k^2$ is the transverse projector (the fluid flow is considered as incompressible), g_0 is a relevant coupling constant (another expansion parameter in the perturbation theory), $1/m$ is a turbulent integral scale Λ' which can be chosen as $\Lambda' = \Lambda$; $0 < \epsilon < 2$ is another parameter of regular expansion of Green's functions which real ("Kolmogorov's") value is $\epsilon_r = 2/3$. Note, that the velocity correlation (5) as specified

above is inherent for numerous simplified models of turbulence introduced by Obukhov, [11] and Kraichnan, [12], see [14] and references therein.

The object of our interest is $G \equiv \langle\langle G(\varphi, v) \rangle\rangle$, where $G(\varphi, v)$ is the Green's function of the differential equation (1) for φ and $\mathbf{v}$ fixed and the doubled angular brackets denote the operations of functional averaging over the known statistics of φ and $\mathbf{v}$.

The stochastic phase factor $\sin c^{-1}\varphi(\mathbf{x})$ can be expanded in powers of φ ,

$$\sin c^{-1}\varphi(\mathbf{x}) = \sum_{\mathbf{n}=1}^{\infty} \frac{(\mathbf{c}^{-1}\varphi)^{2\mathbf{n}-1}(\mathbf{x})}{2\mathbf{n}-1!}$$

as well as the averaged product of two phase factors is expressed by a series of various statistical momenta,

$$\langle \sin c^{-1}\varphi(\mathbf{x}) \sin \mathbf{c}^{-1}\varphi(\mathbf{y}) \rangle = \sum_{\mathbf{n}, \mathbf{m}=1}^{\infty} \frac{\langle (\mathbf{c}^{-1}\varphi)^{2\mathbf{n}-1}(\mathbf{x}) (\mathbf{c}^{-1}\varphi)^{2\mathbf{m}-1}(\mathbf{y}) \rangle}{2\mathbf{n}-1! 2\mathbf{m}-1!}.$$

In order to sidestep consideration of the onset of the problem we compute first the Green's function of the simplified equation with no terms proportional to $\varphi^k(\mathbf{x})$, $k > 1$. Contributions of the higher statistical momenta then are to be determined by various composite operators $F_k = \langle c^{-k}\varphi^k(\mathbf{x}) \rangle$ which we study in the forthcoming sections.

We conclude this section by an explanation of the physical meaning of propagator G . The Fourier transformed propagator $G(k)$, $k = |\mathbf{k}|$ is to be computed in the framework of diagrammatic technique of Feynman. Assuming that initially the marker was concentrated in the source of frame, $J(\mathbf{x}, t) = \delta(\mathbf{x})f(t)$, one obtains the relevant concentration profile of the traveling wave type, $u(\mathbf{r}, t) = u(r)f(t - r\varepsilon/v)$ by the expression

$$u(r) = (2\pi)^{-d} \int d\mathbf{x} G(k) \exp i\mathbf{k}\mathbf{x}. \quad (6)$$

Another practically important problem is to define a concentration distribution profile in a porous half-space. The flat boundary of the medium plays the role of a source term in the right hand side of the equation (1). In case of the porous half-space $x_1 \equiv r \geq 0$, the concentration profile $\langle u(r, t) \rangle$ inside the porous medium is determined by the equation (1) with a source term of the form $J(\mathbf{x}, t) = \delta(x_1)f(t)$ and yielded by the same formula (6) with $d = 1$.

Notice that (6) allows one to bring about the angular integration for arbitrary value of space dimension d ,

$$u(r) = (2\pi)^{-d/2} r^{1-d/2} \int_0^\infty dk G(k) k^{d/2} J_{d/2-1}(kr). \quad (7)$$

where J_ν is the Bessel function. Qualitatively, the solution of (7) is always an outgoing damping traveling wave of concentration which decay rate is of essentially interest for practical applications. We shall discuss the calculation of this rate below in Section 4.

$$G(\varphi, \mathbf{v}) = \text{---}\text{---}\text{---} + \text{---}\text{---}\text{---} + \text{---}\text{---}\text{---} + \dots \quad (9)$$

The relevant action functional has the form

$$S(\varphi, \mathbf{v}, \psi) = \int \mathbf{dx} d\mathbf{t} \left[-\frac{1}{2} \varphi \mathbf{D}_\varphi^{-1} \varphi - \frac{1}{2} \mathbf{v} \mathbf{D}_\mathbf{v}^{-1} \mathbf{v} + \psi^\dagger (\mathbf{L}_0 \psi + \varphi \partial_t \psi + (\mathbf{v} \partial) \psi) \right] + \text{C.C.} \quad (10)$$

where *C.C.* denotes a complex conjugate part, D_φ and D_v are the correlation functions (3) and (5) consequently. Averaging the product $\psi(\mathbf{x}, t) \psi^\dagger(\mathbf{y}, t)$ over the complete set of fields $\{\varphi, \mathbf{v}, \psi\}$ for the statistical weight $\exp S(\varphi, \mathbf{v}, \psi)$, one computes the dressed propagator (the correlation function G) of the "physical" fields,

$$G(\mathbf{x}, \mathbf{y}) = \langle \langle \psi(\mathbf{x}, t) \psi^\dagger(\mathbf{y}, t) \rangle \rangle = C \int D\varphi D\mathbf{v} D\psi \psi(\mathbf{x}, \mathbf{t}) \psi^\dagger(\mathbf{y}, \mathbf{t}) \exp \mathbf{S}(\varphi, \mathbf{v}, \psi). \quad (11)$$

in which the factor C is defined by the normalization condition $\langle \langle 1 \rangle \rangle = 1$.

The correspondence between the original problem (1) and the field theory (10) is not exact, since in the field propagator (11) there are redundant diagrams containing closed loops of the $\psi\psi^\dagger$ -lines. These diagrams were contributed to the correlation functions $\langle \varphi\varphi \rangle$ and $\langle v_i v_j \rangle$ which we think of to be exact as defined in (3) and (5). In principle, these diagram are to be eliminated; one can simply say that the original problem is equivalent to the field theory (10) except the diagrams containing the closed loops of lines of "physical" fields $\psi\psi^\dagger$. This elimination does not affect the theory, since, first, we are not going to perform any functional variables transformations, which could mix up the redundant diagrams with those we need, and second, the renormalization group perfectly well permits such an elimination of a certain class of diagrams.

We also note that all closed loops of $\psi\psi^\dagger$ -lines with no time derivatives in a cycle are automatically equal to zero in the theory (10), since it contains retarded and advanced functions. In particular, from the fact of elimination of all diagrams containing closed loops of oriented lines it follows that

- (i) correlation functions of any number $n > 2$ of the stochastic field φ are trivial,

$$\langle \varphi(\mathbf{x}_1) \dots \varphi(\mathbf{x}_n) \rangle = \mathbf{0};$$

- (ii) correlation functions of any number $n > 2$ of the stochastic field $\mathbf{v}(\mathbf{x}, \mathbf{t})$ are trivial,

$$\langle \mathbf{v}(\mathbf{x}_1, \mathbf{t}) \dots \mathbf{v}(\mathbf{x}_n, \mathbf{t}) \rangle = \mathbf{0};$$

(iii) any mixed correlation functions of stochastic fields $\varphi(\mathbf{x})$ and $\mathbf{v}(\mathbf{x}, \mathbf{t})$ are trivial,

$$\langle \varphi(\mathbf{x}_1) \dots \mathbf{v}(\mathbf{x}_n, \mathbf{t}) \rangle = \mathbf{0}.$$

Furthermore, in the model (10), the odd multipoint correlation functions of scalar field vanish, while the even single-time functions satisfy linear partial differential equations. The solutions for the pair correlation function can be obtained explicitly in analogy with that of in the Kraichnan model of turbulence [12].

Finally, it is essential that the action (10) is invariant under the following field transformations:

$$\begin{aligned} \psi(\mathbf{x}, t) &\rightarrow \psi(\mathbf{x} + \mathbf{s}, t + \tau), \varphi_b(\mathbf{x}, t) \rightarrow \varphi(\mathbf{x} + \mathbf{s}, t + \tau) - b(x), \\ \mathbf{v}_a(\mathbf{x}, t) &\rightarrow \mathbf{v}(\mathbf{x} + \mathbf{s}, t + \tau) - \mathbf{a}(t), \end{aligned} \quad (12)$$

with two parameters: an arbitrary vector function of time $\mathbf{a}(t)$ decreasing at $t \rightarrow -\infty$, $\mathbf{s}(t) = \int_{-\infty}^t \mathbf{a}(\mathbf{t}') d\mathbf{t}'$ and an arbitrary scalar function $b(\mathbf{x})$ fading out at $|\mathbf{x}| \rightarrow \infty$, $\tau(\mathbf{x}) = \int_{\xi > |\mathbf{x} - \mathbf{x}'| \geq \mathbf{w}} \mathbf{b}(\mathbf{x}') d\mathbf{x}'$.

The integral turbulent scale as well as the dissipative length are taken into account in the model (10) by the parameters of infrared cut off m and ultraviolet cut off Λ . The Ward identities which express the invariance of the model (10) under the transformations (12) guarantee that all UV singularities of diagrams are subtracted out in each order of perturbation theory, so that all correlation functions have finite limits in $l_d \rightarrow 0$ and do not depend on l_d in the inertial range. In the region $m \rightarrow 0$ there are strong infrared singularities (powers of m) in the perturbation theory. However, these singularities are not related to the dynamic interactions of eddies which form the spectra of the propagator G but to the kinematic effect of dragging of small eddies by the large ones [13]. This effect can be eliminated completely by taking of the frame of reference which moves with an arbitrary speed of the large-scale eddies.

Since these singularities do not contribute to the spectrum for G , we will not take them into consideration, assuming that the frame of reference moves with some arbitrary speed v along the direction of flow. By the way, the solutions for concentration is to be a traveling wave with the time argument $t - r\varepsilon/v$, as it was proven in [4].

3.2 Dimensional Counting. Action Functional with a "Time-diffusive" Term Included

We start now with the dimensional analysis of model (10). Since the model (10) is of two scales, one can introduce two independent canonical dimensions to each quantity F in the theory (the momentum dimension, d_F^k , and the frequency dimension, d_F^ω). Then, based

on d_F^k and d_F^ω , on can introduce a total canonical dimension $d_F = d_F^k + 2d_F^\omega$ (in the free theory, $\partial_t \sim \Delta$). Assuming $d_k^k = d_\omega^\omega = 1$, one finds out the dimensions of quantities in (10) (see Tab. 1). Nonrenormalized parameters in the Tab. 1 are supplied by the index "0".

Superficial *ultraviolet* (UV) divergences, whose removal requires counterterms, can present only in those Green's functions Γ , which canonical dimension

$$\delta_\Gamma = d + 2 - d_\phi N_\phi \quad (13)$$

is a nonnegative number, [15]. Here, N_ϕ is the number of corresponding fields entering the function Γ and $\phi = \{\varphi, \mathbf{v}, \psi\}$.

In the model (10) the derivative ∂ at the vertex $\psi(\mathbf{v}\partial)\psi^\dagger$ can be moved onto the field ψ by virtue of the vector field $\mathbf{v}$ is transversal. Similarly, the time derivative ∂_t at the vertex $c^{-1}\psi\varphi\partial_t\psi^\dagger$ can be moved onto the field ψ by virtue of the field φ does not depend on time. It decreases effectively the real index of divergence: $\delta_\Gamma^{eff} = \delta_\Gamma - N_\psi$.

From the dimensions in the Tab. 1, taking into account the auxiliary considerations (i)-(iii) of the previous Section as well as those of discussed above, one can find out that for any d superficial divergences can only exist in the 1-irreducible function $\langle\psi\psi^\dagger\rangle$ for which $\delta_\Gamma = 2$, $\delta_\Gamma^{eff} = 0$. The corresponding counterterms must contain either two symbols ∂ and is therefore reduced to $\psi\Delta\psi^\dagger$ or two time derivatives along with the squared c^{-1} , $\nu c^{-2}\psi\partial_t^2\psi^\dagger$. Inclusion of these counterterms into the action functional (10) cannot be reproduced by a simple multiplicative renormalization because of the latter counterterm is not formally present in the action functional (10).

To ensure the multiplicative renormalizability, we introduce an additional term $\nu_0 c_0^{-2} \times \psi\partial_t^2\psi^\dagger$ into (10) supposing c_0 to be a bare parameter; it leads to a renormalized action functional of the form:

$$S_R(\varphi, \mathbf{v}, \psi) = \int \mathbf{dx} \mathbf{dt} \left[-\frac{1}{2}\varphi \mathbf{D}_\varphi^{-1} \varphi - \frac{1}{2}\mathbf{v} \mathbf{D}_\mathbf{v}^{-1} \mathbf{v} + \psi^\dagger (\partial_t \psi + \mathbf{Z}_1 \nu \Delta \psi + \mathbf{Z}_2 \nu c^{-2} \partial_t^2 \psi + \right. \\ \left. + c^{-1} \varphi \partial_t \psi + (v\partial)\psi) \right] + C.C. \quad (14)$$

The inclusion of the new term proportional to c^{-2} corresponds to adding a term of the type $\nu c^{-2} \partial_t^2 u(\mathbf{x}, \mathbf{t})$ into (1). Notice that the inclusion of the $\nu c^{-2} \partial_t^2 u(\mathbf{x}, \mathbf{t})$ -term (a "time-diffusive" term) into the diffusion equation in the presence of axial depending porosity for the first time was proven in [4] for the case of particular stochastic processes originating the large scale porosity fluctuations. Our approach allows to justify this result of [4] and to generalize it for any stochastic law for a porous structure. The coefficient νc^{-2} is analogous to the time parameter τ which was introduced in [4] to characterize time of the diffusion process measured in the time scale of the carrier fluid.

Taking into account the new term in the diffusion equation (1), one obtains instead of (8) in the limit $(k\xi)^2 \gg 1$

$$\langle \psi(\mathbf{k}, t) \psi^\dagger(-\mathbf{k}, t') \rangle_0 \simeq c e^{-ck|t'-t|} / 2\nu k. \quad (15)$$

The damping of correlations in (15) as $c \rightarrow 0$ ($\xi \rightarrow \infty$) is an essential feature of the discussed approximation.

However, we may investigate the initial theory with no such an addition, if we solve the RG equations obtained in extended model with the initial condition $c_0^{-2} = 0$, *i.e.*, if we suppose that either the diffusion process goes up extremely fast or the correlation length ξ is very short due to high dense packed bed. In both cases the domain of discussed microscopic phenomena happens to be small enough and the fluid flow behavior in the packed bed column may be well investigated within the framework of usual porous medium theory dwelling on the constant averaged porosity.

In presence of the time-diffusive term the solutions (7) exhibit a decay in time which rate is determined by the singularity of propagator G , $k^2 = \chi^2$, where $\chi^2 = \nu\omega^2/c^2$ in Fourier representation is analogous to the squared relative refractive index in optics. In a simple pole approximation for $G^{-1} = k^2 - \chi^2$ the Bessel function in (7) can be replaced by its asymptotics at large kr that yields the large distance asymptotics for $u(r)$:

$$u(r) = \frac{i}{2\chi} \left(\frac{\chi}{2\pi i r} \right)^{(d-1)/2}.$$

In this case $Im\chi$ defines the extinction coefficient $Im\chi \sim 1/l_0$, where l_0 is the damping length.

3.3 Renormalization-Group Equations, Critical Scaling, and Empirical Laws of Kolmogorov and Richardson

The UV-divergences (in our case poles in ϵ and δ in diagrams) of the extended model considered (with a "time-diffusive" term included) are removed by the multiplicative renormalization procedure. It amounts to the following: the initial action $S(\Phi)$ is referred to as nonrenormalized, its parameters and coupling constants are referenced to as bare; these are considered as some functions (remaining to be determined) of new renormalized parameters and coupling constants.

The renormalized action functional (14) is a function of renormalized coupling constants and parameters:

$$\begin{aligned} c_0 &= cZ_c, & g_0 &= gM^\epsilon Z_g, \\ \eta_0 &= \eta M^{2\delta} Z_\eta, & \nu_0 &= \nu Z_\nu, \end{aligned} \quad (16)$$

where all renormalization constants Z_a are the functions of two independent quantities Z_1 and Z_2 :

$$\begin{aligned} Z_\nu &= Z_1, & Z_g &= Z_1^{-1}, \\ Z_\eta &= Z_1^{-5} Z_2^2, & Z_c &= Z_1^{1/2} Z_2^{-1/2}, \end{aligned} \quad (17)$$

which can be calculated within the framework of diagram technique. We choose the simplest form of subtraction scheme where the divergences are presented as the bare poles in regularization parameters ϵ and δ (so called minimal subtraction scheme (MS)); M is the renormalization mass parameter, g , η , and ν are renormalized analogues of the bare parameters g_0 , η_0 , and ν_0 . $Z_a = Z_a(g, \eta, c, \epsilon, \delta, d)$ are the renormalization constants. The relations (16) and (17) result from the absence of renormalization for the nonlocal contributions of D_φ and D_v in the action functional (14), so that $g_0\nu_0 = gM^\epsilon\nu$ and $\eta_0\nu_0^3/c_0^{-2} = M^{2\delta}\eta\nu^3/c^{-2}$. No renormalization of the fields ψ and $\mathbf{v}$ and "masses" are required, *i.e.* $Z_\psi = Z_v = 1$ and $M_0 = M$, $m_0 = m$, $Z_m = Z_M = 1$.

The only field φ requires renormalization, $\varphi = \varphi_R Z_\varphi$, $Z_\varphi = Z_1^{1/2} Z_2^{-1/2}$, so that the renormalized Green's function W^R meet the relation

$$W^R(g, \eta, c, \nu, M) Z_\varphi^{N_\varphi} = W(g_0, \eta_0, c_0, \nu_0). \quad (18)$$

W^R are UV-finite function (they are finite in the limits $\epsilon \rightarrow 0$, $\delta \rightarrow 0$) for fixed parameters a .

The RG equations are written for the functions W^R which differ from the initial W only by normalization and, thus, can be used equally validly for critical scaling analysis. To derive these equation one can note that the requirement of eliminating singularities does not determine the functions $e_0 = e_0(e, \epsilon, \delta)$, $e = \{g, \eta, c, \nu\}$, uniquely because of the value of M is not fixed by any physical condition. Variation of M for fixed values of bare parameters e_0 leads to variations of e and Z_e , Z_φ . Following the standard notation, we denote by D_M the differential operator $M\partial_M$ for fixed e_0 . Applying it on both sides of (18) leads to the basic RG equation

$$[D_M + \beta_g \partial_g + \beta_c \partial_c + \beta_\eta \partial_\eta - \gamma_\nu D_\nu + n_\varphi \gamma_\varphi] W^R = 0, \quad (19)$$

where we have used $D_x \equiv x\partial_x$ for any parameters of the renormalization theory; for any Z_i

$$\gamma_i \equiv D_M \ln Z_i, \quad \beta_\alpha \equiv D_M \alpha, \quad \alpha \equiv \{g, \eta, c\}. \quad (20)$$

These identities determine the β -functions of the theory considered,

$$\begin{aligned} \beta_g &= -g(\epsilon - \gamma_1), \\ \beta_\eta &= -\eta(2\delta - 5\gamma_1 + 2\gamma_2), \\ \beta_c &= -\frac{\epsilon}{2}(\gamma_1 - \gamma_2). \end{aligned} \quad (21)$$

and the anomalous dimensionalities γ_i . One calculates the renormalization constants Z_1 and Z_2 from the diagrams of perturbation theory (for example, up to one-loop order from the diagrams depicted in (9)) and then γ_i , β_α functions. All γ - and β - functions are constructed as series in g , η , c , and the functions γ_i depend on neither ϵ nor δ .

Actually, the only reason for calculation of renormalization constants in the theory considered is to prove the existence of *infrared* (IR)-attractive fixed points $\{g_*, \eta_*, c_*\}$ of the RG equations (19) in the space of renormalized charges such that $\beta_\alpha(g_*, \eta_*, c_*) = 0$ and $\omega_{\alpha\zeta} \equiv \partial_\alpha \beta_\zeta(g_*, \eta_*, c_*) > 0$. The values of $\gamma_i(g_*, \eta_*, c_*)$ then can be found from (21) exactly apart from diagram calculations, which are, in fact, pretty standard.

There is a line of IR-fixed points in the model discussed:

$$g_* = \mathcal{A}\epsilon, \quad \eta_* = \mathcal{A}\delta, \quad \forall c_*, \quad \mathcal{A} = (4\pi)^{d/2} \Gamma(d/2) d(d-1)^{-1}, \quad (22)$$

which exists in the physical region $\{g, \eta, c\} > 0$. Along this line

$$\gamma_{1*} = \epsilon, \quad \gamma_{2*} = 2\delta/3, \quad (23)$$

with no corrections of order ϵ^2 , δ^2 and so on.

There are no new critical exponents in the model considered since the critical dimensionalities of φ , c , and $\mathbf{v}$ are just equal to the Kolmogorov's value $-1/3$ as it defined from the phenomenological assumptions (4) and (5), and the dimensionality of the field ψ is equal to its canonical dimension since it has no anomalous corrections ($\gamma_\psi = 0$),

$$\Delta_\varphi = \Delta_c = \Delta_v = -1/3, \quad \Delta_\psi = d/2, \quad \Delta_{\{\psi^\dagger \psi\}} = d. \quad (24)$$

The results expressed in the latter relation (24) can be applied to the study of relative diffusion, *i.e.*, spreading of an admixture cloud consisting of a large number of particles, [16]. If we could label one of the particles in the cloud at time $t = 0$ in the source of a frame $\mathbf{x} = \mathbf{0}$, then the effective radius of the cloud R in the moment $t > 0$ satisfies the relation

$$R^2 = \int d\mathbf{x} \, \mathbf{x}^2 \langle \psi^\dagger(\mathbf{x}, \mathbf{t}) \psi(\mathbf{0}, \mathbf{0}) \rangle. \quad (25)$$

Taking into account that $\Delta_R \equiv -1$, from (24) and (25) one can obtain that $\Delta[dR^2/dt] = -2 - \Delta_t$, where Δ_t is the Kolmogorov's dimensionality of time $\Delta_t = -2 + \gamma_1 = -2/3$. It leads to $dR^2/dt \sim R^{4/3}$, which was to be shown in the Richardson's Four-Thirds Law, [16].

From the first relation in (24) one can easily derive the well-known Kolmogorov's Law of Five-Thirds, [16].

We should stress that the indices (24) as well as their consequences (Richardson's and Kolmogorov's Laws) are the universal features of the model (14), *i.e.*, they do not depend

on the certain positions of fixed points of the RG-transformation of variables in the domain of IR-stability. Physically, this fact means that these properties do not depend on the parameters of the model $\{g, \eta, c, \nu\}$. However, the scaling function $\mathcal{F}$ for the propagator G (and consequently the amplitude factors in the Kolmogorov's and Richardson's Laws, *i.e.*, the Kolmogorov constant and the effective diffusivity coefficient) do depend on $\{g, \eta, c, \nu\}$. We discuss it in more details later on.

Another important note should be made on the role of parameter c for the static (*i.e.*, of equal time) spectra. As we have mentioned before the parameter c determines the diffusion rate in the microscopic scale by means of porosity fluctuations. However, the results (22) and (24) demonstrate that c does not affect the universal properties of the static quantities of model but c can only change the amplitude factors of empirical laws.

3.4 Solutions of RG-Equations, Asymptotics for the correlation function G and the Extinction Coefficient χ

Now we derive the solutions of RG differential equations (19) which give the IR - asymptotics ($r \gg w$) for G . In renormalized variables the propagator G can be expressed in the form

$$G^{-1} = -k^d R(s, g, \eta, c, y, z), \quad (26)$$

where R is some function of dimensionless arguments

$$s \equiv k/M, \quad y \equiv \omega/\nu M^2, \quad z \equiv \chi^2/M^2, \quad (27)$$

which meets the RG-equation

$$[-D_s + \beta_g \partial_g + \beta_\eta \partial_\eta + \beta_c \partial_c - (2 - \gamma_1)D_y + (2\gamma_1 - \gamma_2)D_z] R = 0. \quad (28)$$

The equation (28) can be solved and the function R can be found out.

The general solution of (28) is an arbitrary function of the first integrals $\bar{e} = \{\bar{g}, \bar{\eta}, \bar{c}, \bar{y}, \bar{z}\}$ which number is one less then the number of arguments of R in (26). They can be founded from the system of equations

$$\frac{ds}{s} = \frac{d\bar{g}}{\beta_g(\bar{g})} = \frac{d\bar{\eta}}{\beta_\eta(\bar{g}, \bar{\eta})} = \frac{d\bar{c}}{\beta_c(\bar{g}, \bar{\eta})} = -\frac{d\bar{y}}{\bar{y}(2 - \gamma_1)} = \frac{d\bar{z}}{\bar{z}(2\gamma_1 - \gamma_2)}, \quad (29)$$

supplied by some normalization conditions for $\bar{e}$. We use the standard ones,

$$\bar{e}_i(s = 1, e_i) = e_i. \quad (30)$$

Traditionally, the first integrals associated with coupling constants are called invariant or running charges. In spite of the model (14) is of three charges the relations between them

are so simple that one can (at least up to one-loop order approximation) solve (29)

$$\ln s = \int_g^{\bar{g}} \frac{dx}{\beta_g(x)} = \int_\eta^{\bar{\eta}} \frac{dx}{\beta_\eta(x)}. \quad (31)$$

These integrations can be brought about explicitly. For invariant charges it, in particular, leads to the expressions

$$\bar{g}(s, g) = \frac{\epsilon g}{\epsilon s^\epsilon + \mathcal{A}g(1 - s^\epsilon)}, \quad \bar{\eta}(s, \eta) = \frac{2\delta\eta}{2\delta s^{2\delta} + 3\mathcal{A}\eta(1 - s^{2\delta})}. \quad (32)$$

Taking into account the normalization condition (30) and the relations (31), for any solution $R(s, g, \eta, c, y, z)$ we have

$$R(s, g, \eta, c, y, z) = \left(\frac{s^\epsilon \bar{g}}{g}\right)^{-2/3} \left(\frac{s^{2\delta} \bar{\eta}}{\eta}\right)^{-2/3} \left(\frac{\bar{c}}{c}\right)^{-2} R(1, \bar{g}, \bar{\eta}, \bar{c}, \bar{y}, \bar{z}). \quad (33)$$

Note, that the scaling function $R(1, \bar{g}, \bar{\eta}, \bar{c}, \bar{y}, \bar{z})$ is not fixed by RG-equations and is usually calculated in the framework of diagram technique.

Under the renormalization group transformations the canonical degrees of freedom are replaced by the scaling ones including anomalies. For the scaling asymptotics $s \rightarrow 0$ of (33) it is expressed in the fixation of the values of invariant charges on their values in the fixed points of RG-transformation (22),

$$\bar{g}(s, g) \rightarrow g_*, \quad \bar{\eta}(s, g, \eta) \rightarrow \eta_*, \quad s \equiv k/M \rightarrow 0. \quad (34)$$

In the practical problem it would rather be convenient to express the asymptotics of correlation function G by virtue of bare parameters e_0 , using the relations of renormalization constants (16) and (17). Finally, one has the asymptotics $s \rightarrow 0$ in the following form

$$G^{-1} \simeq A k^d, \quad (35)$$

where $A = -\left(\frac{g_0}{g_*}\right)^{-2/3} \left(\frac{\eta_0}{\eta_*}\right)^{-2/3} \left(\frac{c_0}{c_*}\right)^{-2} R(1, \bar{g}, \bar{\eta}, \bar{c}, \bar{y}, \bar{z})$.

The essential feature of the model (14) is traced in the dividing out of renormalization constants in the IR-asymptotics of Green's function as well as disappearing of the renormalization mass parameter M , which functions in (35) are commended to the bare parameters g_0, η_0, c_0, ν_0 .

For the extinction index χ from (29) we have

$$\bar{\chi}^2 = \chi^2 \exp \int_g^{\bar{g}} dx \frac{2\gamma_1(x)}{\beta_g(x)} - \chi^2 \exp \int_\eta^{\bar{\eta}} dx' \frac{\gamma_2(x')}{\beta_\eta(x')}. \quad (36)$$

In the analogous manner one can obtain the relevant scaling asymptotics:

$$\chi^2 \simeq_{s \rightarrow 0} k^{8/3} \left(\frac{g_0}{g_*}\right)^2 \left(\frac{\eta_0}{\eta_*}\right)^{-1} \left(\frac{c_0}{c_*}\right)^{-2} \bar{\chi}(1, \bar{g}, \bar{\eta}, \bar{c}), \quad (37)$$

which corresponds to the following damping time-spectrum for the dynamic solutions G ,

$$G(t - t') \sim \exp -a_0 k^{2/3} |t - t'|, \quad (38)$$

where $a_0 = \left(\frac{g_0}{g_*}\right)^2 \left(\frac{\eta_0}{\eta_*}\right)^{-1} \left(\frac{c_0}{c_*}\right)^{-2}$. Using (7), one can compute the asymptotic value of static concentration spectrum (with no time dependence) in the case of strong axial porosity fluctuations in the three dimensional space

$$u(r) \sim \frac{1}{r}. \quad (39)$$

This asymptotics replaces the gaussian one for the region much large then w , so that the ordinary gaussian spectrum for this region is to be replaced by k^{-2} for $r \gg w$, see Fig. 2.

4 Renormalization and Critical Dimensions of Composite Operators $c^{-n}\varphi^n(\mathbf{x})$

We have considered the model (14) which corresponds to an unphysical version of the packed beds problem. Now we are going to generalize the developed technique to the case of nongaussian fluctuative component of the porosity field $\tilde{\varepsilon}$.

In the Section 2 we have explained that all amendments due to deviations from the gaussian distribution for $\tilde{\varepsilon}$ can be, in principle, taken into account (at least they can be well-estimated) as a series of power-like composite operators $\langle c^{-n}\varphi^n(\mathbf{x}) \rangle$ in the framework of renormalization-group approach for the model (14). Now we calculate the precise critical indices for each power-like operator.

The canonical dimensions of operators $\langle c^{-n}\varphi^n(\mathbf{x}) \rangle$ are trivial, $d_F = -nd_c + nd_\varphi = 0$, $\forall n$. From the Tab. 1. and (13) one can see that the only diagrams with insertions of power-like operators which contain the superficial divergences have an arbitrary number of external "tails" of the $c^{-1}\varphi$ -type. However, at least one of such an external "tail" is attached either to the vertex $\psi^\dagger(\mathbf{v}\partial)\psi$ or $\psi^\dagger(c^{-1}\varphi\partial_t)\psi$, so that at least one derivative ∂ or ∂_t appears as an extra factor in the diagram, and, consequently, the real index of divergence is necessarily negative.

This means that all operators discussed require no counterterms, *i.e.*, they are UV-finite. The same result can be readily reproduced by consideration of the Ward identities which express the invariance of the model (14) with respect to the field transformations (12). Generally speaking, the model (14) is invariant under the transformations (12) but the operators $\langle c^{-n}\varphi^n(\mathbf{x}) \rangle$ are not, so that they cannot have counterterms and cannot be UV-divergent. Finally, for the critical indices of power-like operators $\langle \varphi^n(\mathbf{x}) \rangle$ one obtains

the simple recurrent relation

$$\Delta[\varphi^n] = -n/3. \quad (40)$$

Note that this relation was not initially clear, and it is a specific feature of the model (14) and those of similar type, [14]. Analogously, one has $\Delta[c^{-n}\varphi^n(\mathbf{x})] = \mathbf{0}$, $\forall \mathbf{n}$.

The latter result exhibits that no one term of power series for the fluctuative component of porosity field $\tilde{\varepsilon}$ can be omitted from the consideration in the equation (1) since their contributions are of equal importance. Thus, we need to consider the theory with all terms of the power series included in the action functional.

5 On the Solutions for Arbitrary $\tilde{\varepsilon}$

The situation similar to that of the problem considered has place in the problem of chemically active scalar admixture [19] and passive scalar admixture convection [20]. Each term of the power-like series happens to be significant in the region we are interested in, so that we cannot limit our consideration by neither the first term $\sim c^{-1}\varphi$ as we did in the Sec. 3 nor any finite number of terms in the power series for a porosity field function (for example, (2)).

Generally speaking, it would lead to the investigation of the problem of an infinite number of dimensionless coupling constants η_n , $n \rightarrow \infty$, *i.e.*, to the renormalized action functional (which is analogous to (14)) of the form

$$\begin{aligned} S_R(\varphi, \mathbf{v}, \psi) = \int \mathbf{dx} \mathbf{dt} \left[-\frac{1}{2}\varphi \mathbf{D}_\varphi^{-1} \varphi - \frac{1}{2}\mathbf{v} \mathbf{D}_\mathbf{v}^{-1} \mathbf{v} + \psi^\dagger (\partial_t \psi + \mathbf{Z}_1 \nu \Delta \psi + \mathbf{Z}_2 \nu \mathbf{c}^{-2} \partial_t^2 \psi + \right. \\ \left. + c^{-1} \varphi \partial_t \psi + \sum_{n=2}^{\infty} c^{-n} \eta_n \varphi^n \partial_t \psi / n! + (v \partial) \psi) \right] + C.C. \end{aligned} \quad (41)$$

However, in contrast with the problems mentioned any of these new terms, $c^{-n} \eta_n \varphi^n \partial_t \psi / n!$, does not contain any UV-divergences in accordance with the general dimensional analysis of Sec. 3.

This means that all the renormalization analysis of the theory (14) is still also valid for (41). In fact, the coupling constants η_n are not the charges in theory (41). Any of the critical indices of the theory (14) does not change in (41) and no one new critical asymptotics appeared.

The basic results (35), (37), (38), (40) are still secure for any power series $c^{-n}\varphi^n$ in (41) *i.e.*, they are valid for any porosity function distributed in the microscopical range.

The only difference between the various models of porosity for the certain critical asymptotics has place in the amplitude factors of Green's functions. They do depend from the certain initial values of the physical parameters of the problem. We believe

that it is the cause of a large dispersion of the various experimental measurements in the problem of the flow through the irregular packed beds.

6 Conclusion

The final conclusion is that the RG method applied to the problem of packed beds with the diffusion coefficient which can arbitrary depend from the concentration of marker in fluid flow demonstrates the existence of a scaling regime in the microscopical range where porosity is subject to the strong fluctuations. The relevant critical indices meet the empirical laws of Kolmogorov and Richardson for any porosity values. However, the nonuniversal quantities of the theory, *i.e.*, the amplitudes of empirical laws are depended on the certain initial values of the parameters as well as the porosity fluctuations. The fluctuative porosity forms the damping time-spectrum for the dynamic correlation functions. Finally, the solution of nonlinear equation of diffusion for the concentration field $u(\mathbf{x}, \mathbf{t})$ is to be a traveling wave with an amplitude damping in space and time. This fact distinguishes the theory considered from the case of ordinary nonlinear diffusion as well as the case of turbulent mixing of passive advection.

We justified the result of [4] on the inclusion of a "time- diffusive" term in the equation (1) for accounting of porosity fluctuation and justified it for an arbitrary stochastic process originating the porosity fluctuations.

7 Acknowledgment

We thank G. Erochenkova for fruitful discussions.

References

- [1] Taylor G.I., Proc. Roy. Soc. London , A**20** (1921), 196
- [2] Legentilhomme P., Legrand J., Comiti J. Journal of applied electrochemistry, **19**, (1989), 263
- [3] Obukhov A.M., in *Atmospheric diffusion and air pollution*, IL, 1962, 138.
- [4] G. Erochenkova , R. Lima Preprint CPT-9/P.3515, July 1997
- [5] R.A. Greenkorn *Flow Phenomena in Porous Media*, Marcel Dekker, Inc. 1983

- [6] Bear, J. *Dynamics of Fluids in Porous Media*, Elsevier, NY (1972)
- [7] Whitaker, S. in *Flow through Porous Media*, American Chemical Society, Washington, D.C., Chap. 2, 31
- [8] Zinn-Justin, J. *Quantum Field Theory and Critical Phenomena*, Clarendon, Oxford, 1990
- [9] Adzhemyan L.Ts., Vasil'ev A.N., Pis'mak Yu. M. Theor. and Math. Phys. **68**, 2, (1987) 770
- [10] A. Munster, *Statistical Thermodynamics*, v. **1**, Springer, 1969
- [11] Obukhov A.M., Izv. Akad. Nauk. SSSR, Geogr. Geophys, **13**, (1949) 58
- [12] R. H. Kraichnan, Phys. Fluids **11**, (1968) 945
- [13] Antonov N.V., JTEP, **105**, 3, 614, (1994)
- [14] Adjemyan L.Ts., Vasil'ev A.N., Antonov N.V., Phys. Rev. E, **58**, 2 (1998), 1823,
- [15] J. Collins, *Renormalization* (Cambridge, 1992)
- [16] A. S. Monin, A. M. Yaglom, *Statistical Fluid Mechanics* (MIT Press, Cambridge, Mass., 1975), Vol. 2.
- [17] Vasil'ev A.N., *Functional Methods in Quantum Field Theory and Statistical Physics*, Gordon and Breach, 1998, ISBN 90-5699-035-7
- [18] L. Ts. Adzhemyan, A.N. Vasil'ev, and Yu. M. Pis'mak, Theor. Math. Phys. **57**, 1131 (1983)
- [19] M. Hnatich, Theor. Math. Phys. **83**, 374 (1990)
- [20] N. V. Antonov, JTEP, **112**, 5 (11), 1649 (1997)

Table 1. Canonical Dimensions of Quantities of the Theory							
F	v, c, φ	ψ	g_0	η_0	g, η	m, μ, M	ν, ν_0
d_F^k	-1	$d/2$	ϵ	2δ	0	1	-2
d_F^ω	1	0	0	0	0	0	1
d_F	1	$d/2$	ϵ	2δ	0	1	0

CAPTION FOR FIGURES

FIGURE 1. On the definition of porosity.

The porosity is directly related to the size of the pores relative to the matrix. If P be a point inside a porous medium on the column axis surrounded by a layer of volume $V(l_i)$, one can define a ratio, $\varepsilon_i = \frac{V_v(l_i)}{V(l_i)}$ in which $V_v(l_i)$ is the volume of void space within volume $V(l_i)$.

FIGURE 2. The deviation of dimensionless concentration spectrum from gaussian for $r \gg w$.

Strongly developed fluctuations of the porosity and velocity fields lead to anomalously small fading off of concentration on the distances $r \gg w$ larger than the size of a pore.